


TNFRSF10A polyclonal antibody

Catalog # PAB30285 Size 100 uL

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human small intestine with TNFRSF10A polyclonal antibody (Cat # PAB30285) shows moderate cytoplasmic positivity in glandular cells.

Specification

Product Description	Rabbit polyclonal antibody raised against partial recombinant human TNFRSF10A.
Immunogen	Recombinant protein corresponding to amino acids 105-139 of human TNFRSF10A.
Sequence	PSSAATIKLHDQSIGTQQWEHSPLGELCPPGSHRS
Host	Rabbit
Reactivity	Human
Form	Liquid
Purification	Antigen affinity purification
Isotype	IgG
Recommend Usage	Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1:50 - 1:200) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH 7.2 (40% glycerol, 0.02% sodium azide).

Storage Instruction

Store at 4°C for short term storage. For long term storage store at -20°C.
Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human small intestine with TNFRSF10A polyclonal antibody (Cat # PAB30285) shows moderate cytoplasmic positivity in glandular cells.

Gene Info — TNFRSF10A

Entrez GeneID [8797](#)

Protein Accession# [O00220](#)

Gene Name TNFRSF10A

Gene Alias APO2, CD261, DR4, MGC9365, TRAILR-1, TRAILR1

Gene Description tumor necrosis factor receptor superfamily, member 10a

Omim ID [603611](#)

Gene Ontology [Hyperlink](#)

Gene Summary The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is activated by tumor necrosis factor-related apoptosis inducing ligand (TNFSF10/TRAIL), and thus transduces cell death signal and induces cell apoptosis. Studies with FADD-deficient mice suggested that FADD, a death domain containing adaptor protein, is required for the apoptosis mediated by this protein. [provided by RefSeq]

Other Designations TNF-related apoptosis inducing ligand receptor 1|cytotoxic TRAIL receptor|death receptor 4

Pathway

- [Apoptosis](#)
- [Cytokine-cytokine receptor interaction](#)
- [Natural killer cell mediated cytotoxicity](#)

Disease

- [Adenocarcinoma](#)
- [Asthma](#)
- [Atherosclerosis](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Colorectal Neoplasms](#)
- [Diabetes Mellitus](#)
- [Disease Models](#)
- [Edema](#)
- [Esophageal Neoplasms](#)
- [Genetic Predisposition to Disease](#)
- [Hematologic Diseases](#)
- [Hodgkin Disease](#)
- [Lung Neoplasms](#)
- [Lymphatic Metastasis](#)
- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Multiple Myeloma](#)
- [Narcolepsy](#)
- [Neoplasm Invasiveness](#)
- [Neoplasm Metastasis](#)
- [Neoplasms](#)

- [Occupational Diseases](#)
- [Ovarian cancer](#)
- [Ovarian Neoplasms](#)
- [Prostate cancer](#)
- [Prostatic Neoplasms](#)
- [Pulmonary Disease](#)
- [Stomach Neoplasms](#)
- [Urinary Bladder Neoplasms](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)