


# CCNE1 polyclonal antibody


Catalog # PAB30542      Size 100 uL

## Applications



### Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human testis with CCNE1 polyclonal antibody (Cat # PAB30542) shows nuclear positivity in cells in seminiferous ducts.



### Immunofluorescence

Immunofluorescent staining of human cell line U-2 OS with CCNE1 polyclonal antibody (Cat # PAB30542) shows positivity in nucleus but excluded from the nucleoli. Antibody staining is shown in green.

## Specification

Product Description	Rabbit polyclonal antibody raised against recombinant human CCNE1.
Immunogen	Recombinant protein corresponding to human CCNE1.
Sequence	RDTMKEDGGAEFSSARSRKRKANVTVFLQDPDEEMAKIDRTARDQCGSQPWDNNAVCADPCLIPTPD

Host	Rabbit
Reactivity	Human
Form	Liquid
Purification	Antigen affinity purification
Isotype	IgG
Recommend Usage	Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1:1000-1:2500) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH 7.2 (40% glycerol, 0.02% sodium azide)
Storage Instruction	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

## Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human testis with CCNE1 polyclonal antibody (Cat # PAB30542) shows nuclear positivity in cells in seminiferous ducts.

- Immunofluorescence

Immunofluorescent staining of human cell line U-2 OS with CCNE1 polyclonal antibody (Cat # PAB30542) shows positivity in nucleus but excluded from the nucleoli. Antibody staining is shown in green.

## Gene Info — CCNE1

Entrez GeneID	<a href="#">898</a>
Protein Accession#	<a href="#">P24864</a>
Gene Name	CCNE1
Gene Alias	CCNE
Gene Description	cyclin E1
Omim ID	<a href="#">123837</a>
Gene Ontology	<a href="#">Hyperlink</a>

**Gene Summary**

The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK2, whose activity is required for cell cycle G1/S transition. This protein accumulates at the G1-S phase boundary and is degraded as cells progress through S phase. Overexpression of this gene has been observed in many tumors, which results in chromosome instability, and thus may contribute to tumorigenesis. This protein was found to associate with, and be involved in, the phosphorylation of NPAT protein (nuclear protein mapped to the ATM locus), which participates in cell-cycle regulated histone gene expression and plays a critical role in promoting cell-cycle progression in the absence of pRB. Two alternatively spliced transcript variants of this gene, which encode distinct isoforms, have been described. Two additional splice variants were reported but detailed nucleotide sequence information is not yet available. [provided by RefSeq]

**Other Designations**

cyclin Es|cyclin Et

**Pathway**

- [Cell cycle](#)
- [p53 signaling pathway](#)
- [Pathways in cancer](#)
- [Prostate cancer](#)
- [Small cell lung cancer](#)

**Disease**

- [Adenocarcinoma](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Disease Progression](#)
- [Esophageal Neoplasms](#)
- [Genetic Predisposition to Disease](#)
- [Neoplasm Invasiveness](#)
- [Neoplasms](#)

- [Ovarian cancer](#)
- [Ovarian Neoplasms](#)
- [Urinary Bladder Neoplasms](#)