


ALDH1A1 polyclonal antibody


Catalog # PAB6018 Size 100 ug

Applications



Western Blot (Tissue lysate)

ALDH1A1 polyclonal antibody (Cat # PAB6018) staining (0.03 ug/mL) of human liver lysate (RIPA buffer, 35 ug total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.



Western Blot (Transfected lysate)

HEK293 overexpressing ALDH1A1 and probed with ALDH1A1 polyclonal antibody (Cat # PAB6018) (mock transfection in first lane), tested by Origene.

Specification	
Product Description	Goat polyclonal antibody raised against synthetic peptide of ALDH1A1.
Immunogen	A synthetic peptide corresponding to C-terminus of human ALDH1A1.
Sequence	C-EVKTVTVKISQKNS
Host	Goat
Theoretical MW (kDa)	54.9
Reactivity	Human, Mouse, Rat



Product Information

Specificity	This antibody may cross-react with ALDH1A2 (GenelD 8854) with one residue difference from this d esign and with ALDH2 (GenelD 217) with two residues difference from this design.
Form	Liquid
Purification	Antigen affinity purification
Concentration	0.5 mg/mL
Recommend Usage	ELISA (1:16000) Western Blot (0.03-0.1 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In Tris saline, pH 7.3 (0.5% BSA, 0.02% sodium azide)
Storage Instruction	Store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

Western Blot (Tissue lysate)

ALDH1A1 polyclonal antibody (Cat # PAB6018) staining (0.03 ug/mL) of human liver lysate (RIPA buffer, 35 ug total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.

Western Blot (Transfected lysate)

HEK293 overexpressing ALDH1A1 and probed with ALDH1A1 polyclonal antibody (Cat # PAB6018) (mock transfection in first lane), tested by Origene.

- Immunohistochemistry
- Enzyme-linked Immunoabsorbent Assay

Gene	Info —	ALDH1A1
		/ \ LUI \

Entrez GenelD	<u>216</u>
Protein Accession#	NP_000680.2
Gene Name	ALDH1A1
Gene Alias	ALDC, ALDH-E1, ALDH11, MGC2318, PUMB1, RALDH1



Product Information

Gene Description	aldehyde dehydrogenase 1 family, member A1 Hyperlink	
Gene Ontology		
Gene Summary	This protein belongs to the aldehyde dehydrogenases family of proteins. Aldehyde dehydrogenas e is the second enzyme of the major oxidative pathway of alcohol metabolism. Two major liver isof orms of this enzyme, cytosolic and mitochondrial, can be distinguished by their electrophoretic mo bilities, kinetic properties, and subcellular localizations. Most Caucasians have two major isozym es, while approximately 50% of Orientals have only the cytosolic isozyme, missing the mitochondri al isozyme. A remarkably higher frequency of acute alcohol intoxication among Orientals than am ong Caucasians could be related to the absence of the mitochondrial isozyme. This gene encode s a cytosolic isoform, which has a high affinity for aldehydes. [provided by RefSeq	
Other Designations	ALDH class 1 acetaldehyde dehydrogenase 1 aldehyde dehydrogenase 1, soluble aldehyde dehydrogenase 1A1 aldehyde dehydrogenase, liver cytosolic retinal dehydrogenase 1 retinaldehyde dehydrogenase 1	

Publication Reference

Developmental expression of aldehyde dehydrogenase in rat: a comparison of liver and lung development.

Yoon M, Madden MC, Barton HA.

Toxicological Sciences 2005 Nov; 89(2):386.

• Retinoic acid imprints gut-homing specificity on T cells.

lwata M, Hirakiyama A, Eshima Y, Kagechika H, Kato C, Song SY.

Immunity 2004 Oct; 21(4):527.

Application: IF, IHC-Fr, Mouse, Small intestine, Spleen

Pathway

- Metabolic pathways
- Retinol metabolism

Disease

- Alcoholism
- Breast cancer
- Breast Neoplasms



- Carcinoma
- Chromosome Deletion
- Cleft Lip
- Cleft Palate
- Disease Models
- Drug Toxicity
- Genetic Predisposition to Disease
- Kidney Failure
- Lymphoma
- Neoplasms
- Ovarian cancer
- Ovarian Neoplasms
- Schizophrenia
- Tobacco Use Disorder