

BCOR polyclonal antibody

Catalog # PAB6440 Size 100 ug

Specification	
Product Description	Goat polyclonal antibody raised against synthetic peptide of BCOR.
Immunogen	A synthetic peptide corresponding to human BCOR.
Sequence	LSATPLYGNVHSW-C
Host	Goat
Theoretical MW (kDa)	188, 186, 192
Specificity	This antibody is expected to recognize both reported isoforms (represented by NP_060215 and NP _065977).
Form	Liquid
Purification	Antigen affinity purification
Concentration	0.5 mg/mL
Quality Control Testing	Antibody Reactive Against Synthetic Peptide.
Recommend Usage	ELISA (1:2000) The optimal working dilution should be determined by the end user.
Storage Buffer	In Tris saline, pH 7.3 (0.5% BSA, 0.02% sodium azide)
Storage Instruction	Store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

ChIP



Enzyme-linked Immunoabsorbent Assay

Gene Info — BCOR	
Entrez GenelD	<u>54880</u>
Protein Accession#	NP_060215;NP_065977
Gene Name	BCOR
Gene Alias	ANOP2, FLJ20285, FLJ38041, KIAA1575, MAA2, MCOPS2, MGC131961, MGC71031
Gene Description	BCL6 co-repressor
Omim ID	<u>300166</u> <u>300485</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	The protein encoded by this gene was identified as an interacting corepressor of BCL6, a POZ/zi nc finger transcription repressor that is required for germinal center formation and may influence a poptosis. This protein selectively interacts with the POZ domain of BCL6, but not with eight other POZ proteins. Specific class I and II histone deacetylases (HDACs) have been shown to interact with this protein, which suggests a possible link between the two classes of HDACs. Several tran script variants encoding different isoforms have been found for this gene. [provided by RefSeq
Other Designations	5830466J11Rik 8430401K06Rik BCL-6 interacting corepressor OTTHUMP00000025766 OTTH UMP00000025768

Publication Reference

The forkhead transcription factor AFX activates apoptosis by induction of the BCL-6 transcriptional repressor.

Tang TT, Dowbenko D, Jackson A, Toney L, Lewin DA, Dent AL, Lasky LA.

The Journal of Biological Chemistry 2002 Apr; 277(16):14255.