

# Goat Anti-Armenian hamster IgG (H&L) secondary antibody

Catalog # PAB9133

Size 2 mg

## Specification

|                            |  |
|----------------------------|--|
| <b>Product Description</b> | Goat anti-Armenian hamster IgG recognizes Armenian hamster IgG whole molecule. This secondary antibody was purified using antigen affinity chromatography.         |
| <b>Immunogen</b>           | Armenian hamster IgG whole molecule  |
| <b>Host</b>                | Goat   |
| <b>Reactivity</b>          | Hamster  |
| <b>Form</b>                | Liquid   |
| <b>Conjugation</b>         | Unconjugated   |
| <b>Purification</b>        | This product was prepared from monospecific antiserum by immunoaffinity chromatography, followed by solid phase adsorption(s) to remove any unwanted reactivities. |
| <b>Recommend Usage</b>     | The optimal working dilution should be determined by the end user.   |
| <b>Storage Buffer</b>      | In 20 mM KH <sub>2</sub> PO <sub>4</sub> , 150 mM NaCl, pH 7.2 (0.01% sodium azide)  |
| <b>Storage Instruction</b> | Store at 4°C. For long term storage store at -20°C.<br>Aliquot to avoid repeated freezing and thawing.   |
| <b>Note</b>                | This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.   |

## Applications

- Western Blot
- Immunohistochemistry
- Enzyme-linked Immunoabsorbent Assay

## Publication Reference

- [Propofol ameliorates endotoxin-induced myocardial cell injury by inhibiting inflammation and apoptosis via the PPAR \$\gamma\$ /HMGB1/NLRP3 axis.](#)

Hui Zhao, Ying Gu, Hai Chen.

Molecular MedicineReports 2021 Mar; 23(3):176.

Application: WB-Ce, WB-Tr, Rat, Rat primary cardiomyocytes