


RecomAb™

PTPRC recombinant monoclonal antibody, clone PTPRC/1783R + PTPRC/1975R


Catalog # RAB00614 Size 100 ug

Applications



Western Blot (Tissue lysate)

Western Blot (tissue lysate) analysis of human spleen.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of human tonsil.

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against full length human PTPRC.
Antibody Species	Rabbit
Immunogen	Recombinant protein corresponding to full-length human PTPRC.
Reactivity	Human
Form	Liquid

Purification	Protein A/G purification
Isotype	IgG
Recommend Usage	Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1-2 ug/mL) Western Blot (1-2 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In 1 mg/mL PBS
Storage Instruction	Store at -20 to -80°C.

Applications

- Western Blot (Tissue lysate)
Western Blot (tissue lysate) analysis of human spleen.
- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)
Immunohistochemical staining of human tonsil.

Gene Info — PTPRC

Entrez GeneID	5788
Protein Accession#	P08575
Gene Name	PTPRC
Gene Alias	B220, CD45, CD45R, GP180, LCA, LY5, T200
Gene Description	protein tyrosine phosphatase, receptor type, C
Omim ID	126200 151460 609532
Gene Ontology	Hyperlink

Gene Summary

The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP contains an extracellular domain, a single transmembrane segment and two tandem intracytoplasmic catalytic domains, and thus belongs to receptor type PTP. This gene is specifically expressed in hematopoietic cells. This PTP has been shown to be an essential regulator of T- and B-cell antigen receptor signaling. It functions through either direct interaction with components of the antigen receptor complexes, or by activating various Src family kinases required for the antigen receptor signaling. This PTP also suppresses JAK kinases, and thus functions as a regulator of cytokine receptor signaling. Four alternatively spliced transcripts variants of this gene, which encode distinct isoforms, have been reported. [provided by RefSeq]

Other Designations

CD45 antigen|T200 glycoprotein|T200 leukocyte common antigen|glycoprotein|leukocyte-common antigen|protein tyrosine phosphatase, receptor type, c polypeptide

Pathway

- [Cell adhesion molecules \(CAMs\)](#)
- [Fc gamma R-mediated phagocytosis](#)
- [Primary immunodeficiency](#)
- [T cell receptor signaling pathway](#)

Disease

- [Arthritis](#)
- [Ascariasis](#)
- [Autoimmune Diseases](#)
- [Cardiomyopathy](#)
- [Diabetes Mellitus](#)
- [Genetic Predisposition to Disease](#)
- [Graves Disease](#)
- [Hashimoto Disease](#)
- [Hepatitis](#)
- [Hepatitis B](#)

- [Hepatitis C](#)
- [HIV Infections](#)
- [Inflammatory Bowel Diseases](#)
- [Lupus Erythematosus](#)
- [Lymphopenia](#)
- [Multiple Sclerosis](#)
- [Paraparesis](#)
- [Scleroderma](#)
- [Severe combined immunodeficiency](#)
- [Thyroiditis](#)