


RecomAb™

CASP1 recombinant monoclonal antibody, clone R05-7H1

Catalog # RAB01539 Size 100 uL

Applications



Western Blot

Western blot analysis of Caspase-1 in HeLa lysates using human Caspase-1 recombinant monoclonal antibody, clone R05-7H1 (Cat # RAB01539).

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human Caspase-1.
Antibody Species	Rabbit
Immunogen	Original antibody is raised against recombinant protein corresponding to human Caspase-1
Theoretical MW (kDa)	Calculated MW: 45 kD
Reactivity	Human
Form	Liquid
Purification	Affinity purification
Isotype	IgG
Recommend Usage	Immunoprecipitation (1:20) Western Blot (1:500-1:1000) The optimal working dilution should be determined by the end user.
Storage Buffer	In 50 mM Tris-Glycine, pH 7.4 (0.15 M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA)

Storage Instruction

Store at 4°C for short term. For long term storage store at -20°C.
Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot

Western blot analysis of Caspase-1 in Hela lysates using human Caspase-1 recombinant monoclonal antibody, clone R05-7H1 (Cat # RAB01539).

- Immunoprecipitation

Gene Info — CASP1

Entrez GeneID[834](#)**Protein Accession#**[P29466](#)**Gene Name**

CASP1

Gene Alias

ICE, IL1BC, P45

Gene Description

caspase 1, apoptosis-related cysteine peptidase (interleukin 1, beta, convertase)

Omim ID[147678](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

This gene encodes a protein which is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce 2 subunits, large and small, that dimerize to form the active enzyme. This gene was identified by its ability to proteolytically cleave and activate the inactive precursor of interleukin-1, a cytokine involved in the processes such as inflammation, septic shock, and wound healing. This gene has been shown to induce cell apoptosis and may function in various developmental stages. Studies of a similar gene in mouse suggest a role in the pathogenesis of Huntington disease. Alternative splicing of this gene results in five transcript variants encoding distinct isoforms. [provided by RefSeq]

Other Designations

CASP1 nirs variant 1|IL1B-convertase|caspase 1|caspase 1, apoptosis-related cysteine protease (interleukin 1, beta, convertase)|interleukin 1-B converting enzyme|interleukin 1-beta convertase

Pathway

- [Amyotrophic lateral sclerosis \(ALS\)](#)

Disease

- [Adenocarcinoma](#)
- [Alzheimer Disease](#)
- [Asthma](#)
- [Bronchiolitis](#)
- [Cardiovascular Diseases](#)
- [Cognition](#)
- [Coronary Artery Disease](#)
- [Diabetes Mellitus](#)
- [Disease Susceptibility](#)
- [Edema](#)
- [Gastrointestinal Stromal Tumors](#)
- [Genetic Predisposition to Disease](#)
- [Hematologic Diseases](#)
- [Hodgkin Disease](#)
- [Infant](#)
- [Inflammation](#)
- [Leukemia](#)
- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Memory](#)

- [Myocardial Infarction](#)
- [Neovascularization](#)
- [Neuropsychological Tests](#)
- [Occupational Diseases](#)
- [Paratyphoid Fever](#)
- [Prostatic Neoplasms](#)
- [Respiratory Syncytial Virus Infections](#)
- [Typhoid Fever](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)