


RecomAb™

F3 recombinant monoclonal antibody, clone R04-7G2

Catalog # RAB01809 Size 100 uL

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin/PFA-fixed paraffin-embedded sections) of human lung cancer with F3 recombinant monoclonal antibody, clone R04-7G2 (Cat # RAB01809). High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human F3.
Antibody Species	Rabbit
Immunogen	Original antibody is raised against a synthetic peptide corresponding to human F3.
Theoretical MW (kDa)	Calculated MW: 33 kD
Reactivity	Human, Mouse, Rat
Form	Liquid
Purification	Affinity purification
Isotype	IgG
Recommend Usage	Immunohistochemistry (1:50-1/100) Western Blot (1:500-1:1000) The optimal working dilution should be determined by the end user.
Storage Buffer	In 50 mM Tris-Glycine, pH 7.4 (0.15 M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA)

Storage Instruction

Store at -20 °C.
Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot
- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin/PFA-fixed paraffin-embedded sections) of human lung cancer with F3 recombinant monoclonal antibody, clone R04-7G2 (Cat # RAB01809). High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Gene Info — F3

Entrez GeneID[2152](#)**Protein Accession#**[P13726](#)**Gene Name**

F3

Gene Alias

CD142, TF, TFA

Gene Description

coagulation factor III (thromboplastin, tissue factor)

Omim ID[134390](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

This gene encodes coagulation factor III which is a cell surface glycoprotein. This factor enables cells to initiate the blood coagulation cascades, and it functions as the high-affinity receptor for the coagulation factor VII. The resulting complex provides a catalytic event that is responsible for initiation of the coagulation protease cascades by specific limited proteolysis. Unlike the other cofactors of these protease cascades, which circulate as nonfunctional precursors, this factor is a potent initiator that is fully functional when expressed on cell surfaces. There are 3 distinct domains of this factor: extracellular, transmembrane, and cytoplasmic. This protein is the only one in the coagulation pathway for which a congenital deficiency has not been described. [provided by RefSeq]

Other Designations

OTTHUMP00000012426|coagulation factor III|tissue factor

Pathway

- [Complement and coagulation cascades](#)

Disease

- [Acute Disease](#)
- [Alzheimer disease](#)
- [Atherosclerosis](#)
- [Brain Ischemia](#)
- [Cardiovascular Diseases](#)
- [Carotid Stenosis](#)
- [Chorioamnionitis](#)
- [Coronary Thrombosis](#)
- [Diabetes Complications](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Fetal Membranes](#)
- [Genetic Predisposition to Disease](#)
- [Hemorrhage](#)
- [Kidney Failure](#)
- [Metabolic Syndrome X](#)
- [Myocardial Ischemia](#)
- [Neoplasms](#)
- [Obstetric Labor](#)
- [Osteoporosis](#)
- [Postoperative Complications](#)
- [Pre-Eclampsia](#)
- [Premature Birth](#)

- [Renal Insufficiency](#)
- [Stroke](#)
- [Thrombosis](#)
- [Vascular Diseases](#)
- [Werner syndrome](#)