


RecomAb™

CBL recombinant monoclonal antibody, clone CbIY700-E1 (FITC)

Catalog # RAB02840 Size 10

ize 100 Reactions

Applications



Flow Cytometry

Flow cytometric analysis of C6 cells cell treated with imatinib and unstained as negative control (blue) or treated with imatinib (red) or with pervanadate (green) and stained using Phospho-c-Cbl (Tyr700) FITC conjugated antibody CblY700-E1.

Specification	
Product Description	Rabbit recombinant monoclonal antibody raised against human CBL.
Antibody Species	Rabbit
Immunogen	A synthetic phospho-peptide corresponding to residues surrounding Tyr700 of human phospho c-Cbl
Reactivity	Human
Form	Liquid
Purification	Protein A+G
lsotype	Rabbit lgG1k
Conjugation Note	FITC
Recommend Usage	Flow Cytometry The optimal working dilution should be determined by the end user.
Storage Buffer	1X PBS, 0.09% Sodium azide, 0.2% BSA
Storage Instruction	Store at 4°C. Do not freeze.

Copyright © 2023 Abnova Corporation. All Rights Reserved.

😵 Abnova

Product Information

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

Flow Cytometry

Flow cytometric analysis of C6 cells cell treated with imatinib and unstained as negative control (blue) or treated with imatinib (red) or with pervanadate (green) and stained using Phospho-c-Cbl (Tyr700) FITC conjugated antibody CblY700-E1.

Gene Info — CBL	
Entrez GenelD	<u>867</u>
Protein Accession#	<u>P22681</u>
Gene Name	CBL
Gene Alias	C-CBL, CBL2, RNF55
Gene Description	Cas-Br-M (murine) ecotropic retroviral transforming sequence
Omim ID	<u>165360</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	The cbl oncogene was first identified as part of a transforming retrovirus which induces mouse pr e-B and pro-B cell lymphomas. As an adaptor protein for receptor protein-tyrosine kinases, it posi tively regulates receptor protein-tyrosine kinase ubiquitination in a manner dependent upon its var iant SH2 and RING finger domains. Ubiquitination of receptor protein-tyrosine kinases terminates signaling by marking active receptors for degradation. [provided by RefSeq
Other Designations	oncogene CBL2

Pathway

- <u>Chronic myeloid leukemia</u>
- Endocytosis
- ErbB signaling pathway
- Insulin signaling pathway
- Jak-STAT signaling pathway

😵 Abnova

- Pathways in cancer
- <u>T cell receptor signaling pathway</u>
- Ubiquitin mediated proteolysis

Disease

- Diabetes Mellitus
- Disease Progression
- Genetic Predisposition to Disease
- Leukemia
- Tobacco Use Disorder