


RecomAb™

STAT3 recombinant monoclonal antibody, clone Stat3Y705-B12 (PE)

Catalog # RAB02872 Size 100 Reactions

Applications



Flow Cytometry

Flow cytometric analysis of Jurkat cells secondary antibody only negative control (blue) or untreated (red) or treated with IFNα IL4 and pervanadate (green) using Phospho-Stat3 (Tyr705) antibody Stat3Y705 B12-PE.

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human STAT3.
Antibody Species	Rabbit
Immunogen	A synthetic phospho-peptide corresponding to residues surrounding Tyr705 of human phospho Stat3
Reactivity	Human
Form	Liquid
Purification	Protein A+G
Isotype	Rabbit IgG1k
Conjugation Note	PE
Recommend Usage	Flow Cytometry The optimal working dilution should be determined by the end user.

Storage Buffer	1X PBS, 0.09% Sodium azide, 0.2% BSA
Storage Instruction	Store at 4°C. Do not freeze.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Flow Cytometry

Flow cytometric analysis of Jurkat cells secondary antibody only negative control (blue) or untreated (red) or treated with IFN α IL4 and pervanadate (green) using Phospho-Stat3 (Tyr705) antibody Stat3Y705 B12-PE.

Gene Info — STAT3

Entrez GeneID	6774
Protein Accession#	P40763
Gene Name	STAT3
Gene Alias	APRF, FLJ20882, HIES, MGC16063
Gene Description	signal transducer and activator of transcription 3 (acute-phase response factor)
Omim ID	102582
Gene Ontology	Hyperlink

Gene Summary

The protein encoded by this gene is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is activated through phosphorylation in response to various cytokines and growth factors including IFNs, EGF, IL5, IL6, HGF, LIF and BMP2. This protein mediates the expression of a variety of genes in response to cell stimuli, and thus plays a key role in many cellular processes such as cell growth and apoptosis. The small GTPase Rac1 has been shown to bind and regulate the activity of this protein. PIAS3 protein is a specific inhibitor of this protein. Three alternatively spliced transcript variants encoding distinct isoforms have been described. [provided by RefSeq]

Other Designations

DNA-binding protein APRF|acute-phase response factor|signal transducer and activator of transcription 3

Pathway

- [Acute myeloid leukemia](#)
- [Adipocytokine signaling pathway](#)
- [Chemokine signaling pathway](#)
- [Jak-STAT signaling pathway](#)
- [Pancreatic cancer](#)
- [Pathways in cancer](#)

Disease

- [Abortion](#)
- [Asthma](#)
- [Autoimmune Diseases](#)
- [Birth Weight](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Bronchiolitis](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Cleft Lip](#)
- [Cleft Palate](#)
- [Colitis](#)
- [Crohn Disease](#)
- [Depressive Disorder](#)
- [Dermatitis](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Disease Susceptibility](#)

- [DNA Damage](#)
- [Eczema](#)
- [Edema](#)
- [Fatty Liver](#)
- [Genetic Predisposition to Disease](#)
- [Genomic Instability](#)
- [Glioblastoma](#)
- [Glioma](#)
- [Hepatitis C](#)
- [Infant](#)
- [Inflammation](#)
- [Inflammatory Bowel Diseases](#)
- [Insulin Resistance](#)
- [Kidney Neoplasms](#)
- [Leukemia](#)
- [Liver Neoplasms](#)
- [Lung Neoplasms](#)
- [Lymphoma](#)
- [Meningeal Neoplasms](#)
- [Meningioma](#)
- [Multiple Sclerosis](#)
- [Neoplasm Metastasis](#)
- [Obesity](#)
- [Ovarian Neoplasms](#)
- [Pulmonary Disease](#)

- [Rectal Fistula](#)
- [Respiratory Syncytial Virus Infections](#)
- [Spondylitis](#)
- [Thyroid Neoplasms](#)
- [Tobacco Use Disorder](#)
- [Tooth Abnormalities](#)