


RecomAb™

# H2AX recombinant monoclonal antibody, clone HisH2AXS139-1B3 (APC)

Catalog # RAB02970

Size 100 Reactions

## Applications



### Flow Cytometry

Flow cytometric analysis of 293T cells, untreated and unstained as negative control (blue) or untreated and stained (red) or treated with UV and PMA and stained (green) using Phospho-Histone H2A.X (Ser139) antibody HisH2AXS139-1B3 APC conjugate.

## Specification

<b>Product Description</b>	Rabbit recombinant monoclonal antibody raised against human H2AX.
<b>Antibody Species</b>	Rabbit
<b>Immunogen</b>	A synthetic phospho-peptide corresponding to residues surrounding Ser139 of human phospho-histone H2A.X
<b>Reactivity</b>	Human
<b>Form</b>	Liquid
<b>Conjugation</b>	APC
<b>Purification</b>	Protein A purification, Protein G purification
<b>Isotype</b>	IgG
<b>Recommend Usage</b>	Flow Cytometry The optimal working dilution should be determined by the end user.

<b>Storage Buffer</b>	1X PBS, 0.09% Sodium azide, 0.2% BSA
<b>Storage Instruction</b>	Store at 4°C. Do not freeze.
<b>Note</b>	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

## Applications

- Flow Cytometry

Flow cytometric analysis of 293T cells, untreated and unstained as negative control (blue) or untreated and stained (red) or treated with UV and PMA and stained (green) using Phospho-Histone H2A.X (Ser139) antibody HisH2AXS139-1B3 APC conjugate.

## Gene Info — H2AFX

<b>Entrez GeneID</b>	<a href="#">3014</a>
<b>Protein Accession#</b>	<a href="#">P16104</a>
<b>Gene Name</b>	H2AFX
<b>Gene Alias</b>	H2A.X, H2A/X, H2AX
<b>Gene Description</b>	H2A histone family, member X
<b>Omim ID</b>	<a href="#">601772</a>
<b>Gene Ontology</b>	<a href="#">Hyperlink</a>
<b>Gene Summary</b>	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene encodes a member of the histone H2A family, and generates two transcripts through the use of the conserved stem-loop termination motif, and the polyA addition motif. [provided by RefSeq]
<b>Other Designations</b>	H2AX histone

## Pathway

- [Systemic lupus erythematosus](#)

## Disease

- [Azoospermia](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [DNA Damage](#)
- [Genetic Predisposition to Disease](#)
- [Lymphoma](#)
- [Oligospermia](#)
- [Ovarian cancer](#)
- [Prostate cancer](#)
- [Prostatic Neoplasms](#)
- [Urinary Bladder Neoplasms](#)