


RecomAb™

RELA recombinant monoclonal antibody, clone NFkBp65S529-H3 (FITC)

Catalog # RAB03072

Size 100 Reactions

Applications



Flow Cytometry

Flow cytometric analysis of HeLa cells untreated and unstained as negative control (blue) or untreated (red) or treated with TNFa + calyculin A (green) and stained using Phospho-NFkB p65 (Ser529) antibody FITC conjugate, NFkBp65S529-H3.

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human RELA.
Antibody Species	Rabbit
Immunogen	A synthetic phospho-peptide corresponding to residues surrounding Ser529 of human phospho-NFK B p65
Reactivity	Human
Form	Liquid
Conjugation	FITC
Purification	Protein A purification, Protein G purification
Isotype	IgG
Recommend Usage	Flow Cytometry The optimal working dilution should be determined by the end user.

Storage Buffer	In PBS (0.2% BSA, 0.09% Sodium azide)
Storage Instruction	Store at 4°C. Do not freeze.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Flow Cytometry

Flow cytometric analysis of HeLa cells untreated and unstained as negative control (blue) or untreated (red) or treated with TNFα + calyculin A (green) and stained using Phospho-NFκB p65 (Ser529) antibody FITC conjugate, NFκBp65S529-H3.

Gene Info — RELA

Entrez GeneID	5970
Protein Accession#	Q04206
Gene Name	RELA
Gene Alias	MGC131774, NFκB3, p65
Gene Description	v-rel reticuloendotheliosis viral oncogene homolog A (avian)
Omim ID	164014
Gene Ontology	Hyperlink
Gene Summary	NFκB1 (MIM 164011) or NFκB2 (MIM 164012) is bound to REL (MIM 164910), RELA, or RELB (MIM 604758) to form the NFκB complex. The p50 (NFκB1)/p65 (RELA) heterodimer is the most abundant form of NFκB. The NFκB complex is inhibited by I-κappa-B proteins (NFκBIA, MIM 164008 or NFκBIB, MIM 604495), which inactivate NFκB by trapping it in the cytoplasm. Phosphorylation of serine residues on the I-κappa-B proteins by kinases (IKBKA, MIM 600664, or IKBKB, MIM 603258) marks them for destruction via the ubiquitination pathway, thereby allowing activation of the NFκB complex. Activated NFκB complex translocates into the nucleus and binds DNA at kappa-B-binding motifs such as 5-prime GGGRNNYYCC 3-prime or 5-prime HGGARNYYCC 3-prime (where H is A, C, or T; R is an A or G purine; and Y is a C or T pyrimidine).[supplied by OMIM]
Other Designations	nuclear factor of kappa light polypeptide gene enhancer in B-cells 3 v-rel avian reticuloendotheliosis viral oncogene homolog A (nuclear factor of kappa light polypeptide gene enhancer in B-cells 3 (p65)) v-rel reticuloendotheliosis viral oncogene homolog

Pathway

- [Acute myeloid leukemia](#)
- [Adipocytokine signaling pathway](#)
- [Apoptosis](#)
- [B cell receptor signaling pathway](#)
- [Chemokine signaling pathway](#)
- [Chronic myeloid leukemia](#)
- [Epithelial cell signaling in Helicobacter pylori infection](#)
- [MAPK signaling pathway](#)
- [Neurotrophin signaling pathway](#)
- [Pancreatic cancer](#)
- [Pathways in cancer](#)
- [Prostate cancer](#)
- [Small cell lung cancer](#)
- [T cell receptor signaling pathway](#)
- [Toll-like receptor signaling pathway](#)

Disease

- [Arthritis](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Cardiovascular Diseases](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Disease Susceptibility](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)

- [Hematologic Diseases](#)
- [HIV Infections](#)
- [Hodgkin Disease](#)
- [Liver Cirrhosis](#)
- [Lymphoproliferative Disorders](#)
- [Multiple Myeloma](#)
- [Occupational Diseases](#)
- [Testicular Neoplasms](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)