RecomAb™

IL2RA recombinant monoclonal antibody, clone AHT107

Catalog # RAB03579 Size 200 ug

Applications



Immunofluorescence

Immunofluorescent staining of Jurkat cells with IL2RA recombinant monoclonal antibody, clone AHT107 (Cat # RAB03579).
Immunofluorescence analysis of paraformaldehyde fixed Jurkat cells immobilized on Shi-fix™ coverslips and stained with the chimeric r version of RAB03579 at 10 ug/mL followed by Alexa Fluor® 488 secondary antibody (2 ug/mL), showing membrane staining. The nuclear stain is DAPI (blue). The isotype control was stained with anti-Fluorescein antibody followed by staining with Alexa Fluor® 488 secondary antibody.
(A) RAB03579
(B) DAPI
(C) Merged channels
(D) Isotype control

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human IL2RA.
Antibody Species	Rabbit
Immunogen	Original antibody is raised against human T lymphoblasts.
Reactivity	Human
Form	Liquid
Isotype	IgG

Recommend Usage	Blocking ELISA Flow Cytometry Immunofluorescence The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.02% Proclin 300)
Storage Instruction	Store at 4°C for 3 months. For long term storage, aliquot and store at -20°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Immunofluorescence

  Immunofluorescent staining of Jurkat cells with IL2RA recombinant monoclonal antibody, clone AHT107 (Cat # RAB03579). Immunofluorescence analysis of paraformaldehyde fixed Jurkat cells immobilized on Shi-fix™ coverslips and stained with the chimeric r version of RAB03579 at 10 ug/mL followed by Alexa Fluor® 488 secondary antibody (2 ug/mL), showing membrane staining. The nuclear stain is DAPI (blue). The isotype control was stained with anti-Fluorescein antibody followed by staining with Alexa Fluor® 488 secondary antibody.
  (A) RAB03579
  (B) DAPI
  (C) Merged channels
  (D) Isotype control

- Enzyme-linked Immunoabsorbent Assay

- Blocking

- Flow Cytometry

Gene Info — IL2RA

Entrez GeneID	3559
Gene Name	IL2RA
Gene Alias	CD25, IDDM10, IL2R, TCGFR
Gene Description	interleukin 2 receptor, alpha
Omim ID	147730 601942
Gene Ontology	Hyperlink

Gene Summary	The interleukin 2 (IL2) receptor alpha (IL2RA) and beta (IL2RB) chains, together with the common gamma chain (IL2RG), constitute the high-affinity IL2 receptor. Homodimeric alpha chains (IL2RA) result in low-affinity receptor, while homodimeric beta (IL2RB) chains produce a medium-affinity r eceptor. Normally an integral-membrane protein, soluble IL2RA has been isolated and determine d to result from extracellular proteolyisis. Alternately-spliced IL2RA mRNAs have been isolated, b ut the significance of each is presently unknown. [provided by RefSeq
Other Designations	Interleukin-2 receptor|OTTHUMP00000019030|OTTHUMP00000044870|interleukin 2 receptor, al pha chain

Pathway

- Cytokine-cytokine receptor interaction
- Endocytosis
- Hematopoietic cell lineage
- Jak-STAT signaling pathway

Disease

- Acquired Immunodeficiency Syndrome
- Addison Disease
- Alopecia Areata
- Alzheimer Disease
- Arthritis
- Asthma
- Atrial Fibrillation
- Autoimmune Diseases
- Bronchiolitis
- Carcinoma
- Cardiovascular Diseases
- Celiac Disease
- Chorioamnionitis

- Colitis
- Connective Tissue Diseases
- Crohn Disease
- Diabetes Mellitus
- Disease Progression
- Edema
- Fetal Diseases
- Fetal Membranes
- Genetic Predisposition to Disease
- Graves Disease
- Hyperparathyroidism
- Hypersensitivity
- Infant
- Infection
- Inflammation
- Lupus Erythematosus
- Measles
- Melanoma
- Multiple Sclerosis
- Mumps
- Musculoskeletal Diseases
- Obstetric Labor
- Prediabetic State
- Pre-Eclampsia
- Pregnancy Complications

- Premature Birth
- Prostatic Neoplasms
- Respiratory Syncytial Virus Infections
- Rubella
- Skin Diseases
- Thyroid Diseases
- Vasculitis
- Vitiligo