

RecomAb™

PML recombinant monoclonal antibody, clone R02-6K4

Catalog # RAB06504 Size 100 uL

| Specification | |
|----------------------|---|
| Product Description | Rabbit recombinant monoclonal antibody raised against human PML. |
| Antibody Species | Rabbit |
| Immunogen | Original antibody is raised against recombinant protein corresponding to human PML. |
| Theoretical MW (kDa) | Calculated MW: 98 kD |
| Reactivity | Human, Mouse, Rat |
| Form | Liquid |
| Purification | Affinity purification |
| lsotype | lgG |
| Recommend Usage | Flow Cytometry (1:50-1:100) Immunohistochemistry (1:50-1:100) Immunofluorescence(1:50-1:200) Immunoprecipitation(1:20) Western Blot (1:500-1:1000) The optimal working dilution should be determined by the end use. |
| Storage Buffer | In PBS, 150 mM NaCI, pH 7.4 (50% glycerol and 0.02% Sodium azide) |
| Storage Instruction | Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing. |
| Note | This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul |

Applications

Western Blot

- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytometry

Gene Info — PML

| Entrez GenelD | <u>5371</u> |
|--------------------|--|
| Gene Name | PML |
| Gene Alias | MYL, PP8675, RNF71, TRIM19 |
| Gene Description | promyelocytic leukemia |
| Omim ID | <u>102578</u> |
| Gene Ontology | Hyperlink |
| Gene Summary | The protein encoded by this gene is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. This phosphoprotein localizes to nuclear bodies where it functions as a transcription factor and tumor suppressor. Its expression is cell-cycle related and it regulates the p53 response to onc ogenic signals. The gene is often involved in the translocation with the retinoic acid receptor alpha gene associated with acute promyelocytic leukemia (APL). Extensive alternative splicing of this g ene results in several variations of the protein's central and C-terminal regions; all variants encode the same N-terminus. Alternatively spliced transcript variants encoding different isoforms have be en identified. [provided by RefSeq |
| Other Designations | promyelocytic leukemia protein promyelocytic leukemia, inducer of tripartite motif protein TRIM19 |

Pathway

- Acute myeloid leukemia
- Pathways in cancer
- Ubiquitin mediated proteolysis



Disease

- Cerebral Hemorrhage
- Genetic Predisposition to Disease
- Hypertension
- Intracranial Hemorrhages
- Leukemia
- <u>Stroke</u>
- Subarachnoid Hemorrhage
- Tobacco Use Disorder