


RecomAb™

BTK recombinant monoclonal antibody, clone 3E5

Catalog # RAB07514 Size 100 uL

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical analysis of paraffin-embedded human tonsil tissue using BTK recombinant monoclonal antibody, clone 3E5 (Cat # RAB07514) on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.

Specification

Product Description Rabbit recombinant monoclonal antibody raised against human BTK.

Antibody Species Rabbit

Immunogen Original antibody is raised against a synthetic peptide corresponding to human BTK.

Reactivity Human

Form Liquid

Purification Affinity chromatography purification

Isotype IgG

Recommend Usage ELISA
Immunohistochemistry (1:50-1:200)
The optimal working dilution should be determined by the end user.

Storage Buffer In PBS, pH7.4 (150 mM NaCl, 0.02% sodium azide and 50% glycerol)

Storage Instruction

Store at -20°C or -80°C.
Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical analysis of paraffin-embedded human tonsil tissue using BTK recombinant monoclonal antibody, clone 3E5 (Cat # RAB07514) on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.

- Enzyme-linked Immunoabsorbent Assay

Gene Info — BTK

Entrez GeneID[695](#)**Protein Accession#**[Q06187](#)**Gene Name**

BTK

Gene Alias

AGMX1, AT, ATK, BPK, IMD1, MGC126261, MGC126262, PSCTK1, XLA

Gene Description

Bruton agammaglobulinemia tyrosine kinase

Omim ID[300300](#) [307200](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

The protein encoded by this gene plays a crucial role in B-cell development. Mutations in this gene cause X-linked agammaglobulinemia type 1, which is an immunodeficiency characterized by the failure to produce mature B lymphocytes, and associated with a failure of Ig heavy chain rearrangement. [provided by RefSeq]

Other Designations

B-cell progenitor kinase|Bruton tyrosine kinase|OTTHUMP00000023676|OTTHUMP00000063593|agammaglobulinemia tyrosine kinase|dominant-negative kinase-deficient Brutons tyrosine kinase|tyrosine-protein kinase BTK

Pathway

- [B cell receptor signaling pathway](#)
- [Fc epsilon RI signaling pathway](#)
- [Primary immunodeficiency](#)

Disease

- [Agammaglobulinemia](#)
- [Asthma](#)
- [Cardiovascular Diseases](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Genetic Diseases](#)
- [HIV Infections](#)
- [Immunologic Deficiency Syndromes](#)
- [Lymphoproliferative Disorders](#)
- [Myelodysplastic Syndromes](#)
- [Severe Combined Immunodeficiency](#)