


RecomAb™

MET recombinant monoclonal antibody, clone 6E5

Catalog # RAB07538 Size 100 uL

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical analysis of paraffin-embedded human breast cancer with MET recombinant monoclonal antibody, clone 6E5 (Cat # RAB07538) on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human MET.
Antibody Species	Rabbit
Immunogen	Original antibody is raised against a synthetic peptide corresponding to human MET.
Reactivity	Human
Form	Liquid
Purification	Affinity chromatography purification
Isotype	IgG
Recommend Usage	ELISA Immunohistochemistry (1:50-1:200) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH7.4 (150 mM NaCl, 0.02% sodium azide and 50% glycerol)

Storage Instruction

Store at -20°C or -80°C.
Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical analysis of paraffin-embedded human breast cancer with MET recombinant monoclonal antibody, clone 6E5 (Cat # RAB07538) on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.

- Enzyme-linked Immunoabsorbent Assay

Gene Info — MET

Entrez GeneID[4233](#)**Protein Accession#**[P08581](#)**Gene Name**

MET

Gene Alias

AUTS9, HGFR, RCCP2, c-Met

Gene Description

met proto-oncogene (hepatocyte growth factor receptor)

Omim ID[114550](#) [164860](#) [605074](#) [611015](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

The proto-oncogene MET product is the hepatocyte growth factor receptor and encodes tyrosine-kinase activity. The primary single chain precursor protein is post-translationally cleaved to produce the alpha and beta subunits, which are disulfide linked to form the mature receptor. Various mutations in the MET gene are associated with papillary renal carcinoma. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]

Other Designations

HGF receptor|OTTHUMP00000069168|SF receptor|met proto-oncogene|met proto-oncogene tyrosine kinase|oncogene MET|scatter factor receptor

Pathway

- [Adherens junction](#)
- [Axon guidance](#)
- [Colorectal cancer](#)
- [Cytokine-cytokine receptor interaction](#)
- [Endocytosis](#)
- [Epithelial cell signaling in Helicobacter pylori infection](#)
- [Focal adhesion](#)
- [Melanoma](#)
- [Pathways in cancer](#)
- [Renal cell carcinoma](#)

Disease

- [Adenocarcinoma](#)
- [Autistic Disorder](#)
- [Carcinoma](#)
- [Cell Transformation](#)
- [Child Development Disorders](#)
- [Chronic Disease](#)
- [Cognition Disorders](#)
- [Disease Progression](#)
- [Gastrointestinal Diseases](#)
- [Genetic Predisposition to Disease](#)
- [Head and Neck Neoplasms](#)
- [Hyperparathyroidism](#)
- [Lung Neoplasms](#)

- [Myopia](#)
- [Nasal Polyps](#)
- [Neoplasm Metastasis](#)
- [Neoplasms](#)
- [Neuropsychological Tests](#)
- [Pancreatic cancer](#)
- [Pancreatic Neoplasms](#)
- [Pulmonary Disease](#)
- [Refractive Errors](#)
- [Rhinitis](#)
- [Schizophrenia](#)
- [Sinusitis](#)
- [Skin Neoplasms](#)
- [Urinary Bladder Neoplasms](#)
- [Werner syndrome](#)