
RecomAb™

MAPK14 recombinant monoclonal antibody, clone 6H8


Catalog # RAB07541 Size 100 uL

Applications




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical analysis of paraffin-embedded human brain tissue using MAPK14 recombinant monoclonal antibody, clone 6H8 (Cat # RAB07541) on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.



Immunofluorescence

Immunofluorescent staining of Hela Cells with MAPK14 recombinant monoclonal antibody, clone 6H8 (Cat # RAB07541). The cells were fixed in 4% formaldehyde, permeated by 0.2% TritonX-100, and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. Nuclear DNA was labeled in blue with DAPI. The secondary antibody was FITC-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L).



Western Blot

Western blot analysis of Hela whole cell lysate, Jurkat whole cell lysate, MCF-7 whole cell lysate, U-87 whole cell lysate, PC-3 whole cell lysate, L02 whole cell lysate, Rat Heart whole cell lysate with MAPK14 recombinant monoclonal antibody, clone 6H8 (Cat # RAB07541).

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human MAPK14.
Antibody Species	Rabbit
Immunogen	Original antibody is raised against a synthetic peptide corresponding to human MAPK14.
Theoretical MW (kDa)	Calculated MW: 42, 3
Reactivity	Human, Rat
Form	Liquid
Preparation Method	Mammalian cell (HEK293) expression system
Purification	Affinity chromatography purification
Isotype	IgG
Recommend Usage	ELISA Immunofluorescence (1:20-1:200) Immunohistochemistry (1:50-1:200) Western Blot (1:500-1:5000) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH7.4 (150 mM NaCl, 0.02% sodium azide and 50% glycerol)
Storage Instruction	Store at -20°C or -80°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

  Immunohistochemical analysis of paraffin-embedded human brain tissue using MAPK14 recombinant monoclonal antibody, clone 6H8 (Cat # RAB07541) on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.

Copyright © 2023 Abnova Corporation. All Rights Reserved.

- Immunofluorescence

  Immunofluorescent staining of Hela Cells with MAPK14 recombinant monoclonal antibody, clone 6H8 (Cat # RAB07541). The cells were fixed in 4% formaldehyde, permeated by 0.2% TritonX-100, and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. Nuclear DNA was labeled in blue with DAPI. The secondary antibody was FITC-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L).

- Enzyme-linked Immunoabsorbent Assay

- Western Blot

  Western blot analysis of Hela whole cell lysate, Jurkat whole cell lysate, MCF-7 whole cell lysate, U-87 whole cell lysate, PC-3 whole cell lysate, L02 whole cell lysate, Rat Heart whole cell lysate with MAPK14 recombinant monoclonal antibody, clone 6H8 (Cat # RAB07541).

Gene Info — MAPK14

Entrez GeneID	1432
Protein Accession#	Q16539
Gene Name	MAPK14
Gene Alias	CSBP1, CSBP2, CSPB1, EXIP, Mxi2, PRKM14, PRKM15, RK, SAPK2A, p38, p38ALPHA
Gene Description	mitogen-activated protein kinase 14
Omim ID	600289
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a member of the MAP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular pro cesses such as proliferation, differentiation, transcription regulation and development. This kinase is activated by various environmental stresses and proinflammatory cytokines. The activation requ ires its phosphorylation by MAP kinase kinases (MKKs), or its autophosphorylation triggered by t he interaction of MAP3K7IP1/TAB1 protein with this kinase. The substrates of this kinase include transcription regulator ATF2, MEF2C, and MAX, cell cycle regulator CDC25B, and tumor suppres sor p53, which suggest the roles of this kinase in stress related transcription and cell cycle regulat ion, as well as in genotoxic stress response. Four alternatively spliced transcript variants of this g ene encoding distinct isoforms have been reported. [provided by RefSeq
Other Designations	Csaids binding protein|MAP kinase Mxi2|MAX-interacting protein 2|cytokine suppressive anti-infl ammatory drug binding protein|p38 MAP kinase|p38 mitogen activated protein kinase|p38alpha Exip|stress-activated protein kinase 2A

Pathway

Copyright © 2023 Abnova Corporation. All Rights Reserved.

- Amyotrophic lateral sclerosis (ALS)
- Epithelial cell signaling in Helicobacter pylori infection
- Fc epsilon RI signaling pathway
- GnRH signaling pathway
- Leukocyte transendothelial migration
- MAPK signaling pathway
- Neurotrophin signaling pathway
- T cell receptor signaling pathway
- Toll-like receptor signaling pathway
- VEGF signaling pathway

Disease

- Cardiovascular Diseases
- Diabetes Mellitus
- Disease Models
- Edema
- Genetic Predisposition to Disease
- HIV Infections
- Narcolepsy
- Obesity
- Ovarian Failure
- Polycystic Ovary Syndrome
- Puberty
- Schizophrenia

Copyright © 2023 Abnova Corporation. All Rights Reserved.

- Thrombophilia
- Tobacco Use Disorder

Copyright © 2023 Abnova Corporation. All Rights Reserved.