RecomAb™ Sm Size

Histone H4 (acetyl K16) monoclonal antibody, clone RM204

Catalog # RAB09453 Size 25 ug

Applications



Western Blot (Cell lysate)

Western Blot analysis of Lane 1: acid extracts of HeLa cell treated with sodium butyrate and Lane 2: acid extracts of HeLa cell untreated with Histone H4 (acetyl K16) recombinant monoclonal antibody, clone RM204 (Cat # RAB09453) at 0.5 ug/mL working concentration, showed a band of Histone H4 acetylated at Lysine 16.

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against of human histone H4 (acetyl K16).
Antibody Species	Rabbit
Immunogen	Original antibody is raised against a synthetic acetyl peptide corresponding to residues surrounding K16 of human Histone H4.
Sequence	N/A
Specificity	This antibody reacts to Histone H4 acetylated at Lysine 16. No cross reactivity with other acetylated Lysines in Histone H4.
Form	Liquid
Purification	Protein A purification
Isotype	IgG
Recommend Usage	ELISA (0.5 ug/mL-1 ug/mL) Western Blot (0.5 ug/mL-2 ug/mL) The optimal working dilution should be determined by the end user.

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Storage Buffer	In PBS (50% glycerol, 1% BSA, 0.09% sodium azide)
Storage Instruction	Store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

- Enzyme-linked Immunoabsorbent Assay
- Western Blot (Cell lysate)

  Western Blot analysis of Lane 1: acid extracts of HeLa cell treated with sodium butyrate and Lane 2: acid extracts of HeLa cell untreated with Histone H4 (acetyl K16) recombinant monoclonal antibody, clone RM204 (Cat # RAB09453) at 0.5 ug/mL working concentration, showed a band of Histone H4 acetylated at Lysine 16.

Gene Info — HIST4H4

Entrez GeneID	121504
Protein Accession#	P62805
Gene Name	HIST4H4
Gene Alias	H4/p, MGC24116
Gene Description	histone cluster 4, H4
Gene Ontology	Hyperlink
Gene Summary	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chro mosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped aro und a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H 4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H4 family. Transcripts from this gene lack polyA tails; inste ad, they contain a palindromic termination element. [provided by RefSeq
Other Designations	histone 4, H4|histone H4

Pathway

Copyright © 2023 Abnova Corporation. All Rights Reserved.

- Systemic lupus erythematosus

Copyright © 2023 Abnova Corporation. All Rights Reserved.