RecomAb™ Sm Size

Histone H2AX monoclonal antibody, clone RM214

Catalog # RAB09468 Size 25 ug

Applications



Immunocytochemistry

Immunocytochemical staining of HeLa cells with Histone H2AX recombinant monoclonal antibody, clone RM214 (Cat # RAB09468) (Red). Actin filaments have been labeled with fluorescein phalloidin (Green).



Western Blot (Cell lysate)

Western Blot analysis of Lane 1: A375, Lane 2: HEK293, Lane 3: HeLa and Lane 4: SK-MEL-2 whole cell lysate with Histone H2AX recombinant monoclonal antibody, clone RM214 (Cat # RAB09468) at 0.5 ug/mL working concentration, showed endogenous Histone H2AX in A375, HEK293, HeLa and SK-MEL-2 cells.

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against of human histone H2AX.
Antibody Species	Rabbit
Immunogen	Original antibody is raised against a synthetic peptide corresponding to C-terminus region of human Histone H2AX.
Sequence	N/A

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Specificity	This antibody reacts to Histone H2AX protein, independent of post-translational modifications. No cr oss reactivity with other histone proteins. .
Form	Liquid
Purification	Protein A purification
Isotype	IgG
Recommend Usage	ELISA (0.2 ug/mL-1 ug/mL) Immunocytochemistry (1 ug/mL-2 ug/mL) Western Blot (0.5 ug/mL-2 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (50% glycerol, 1% BSA, 0.09% sodium azide)
Storage Instruction	Store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

- Immunocytochemistry

  Immunocytochemical staining of HeLa cells with Histone H2AX recombinant monoclonal antibody, clone RM214 (Cat # RAB09468) (Red). Actin filaments have been labeled with fluorescein phalloidin (Green).

- Enzyme-linked Immunoabsorbent Assay

- Western Blot (Cell lysate)

  Western Blot analysis of Lane 1: A375, Lane 2: HEK293, Lane 3: HeLa and Lane 4: SK-MEL-2 whole cell lysate with Histone H2AX recombinant monoclonal antibody, clone RM214 (Cat # RAB09468) at 0.5 ug/mL working concentration, showed endogenous Histone H2AX in A375, HEK293, HeLa and SK-MEL-2 cells.

Gene Info — H2AFX

Entrez GeneID	3014
Protein Accession#	P16104
Gene Name	H2AFX
Gene Alias	H2A.X, H2A/X, H2AX

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Gene Description	H2A histone family, member X
Omim ID	601772
Gene Ontology	Hyperlink
Gene Summary	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chro mosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, an d H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and f unctions in the compaction of chromatin into higher order structures. This gene encodes a membe r of the histone H2A family, and generates two transcripts through the use of the conserved stem-l oop termination motif, and the polyA addition motif. [provided by RefSeq
Other Designations	H2AX histone

Pathway

- Systemic lupus erythematosus

Disease

- Azoospermia
- Breast cancer
- Breast Neoplasms
- DNA Damage
- Genetic Predisposition to Disease
- Lymphoma
- Oligospermia
- Ovarian cancer
- Prostate cancer
- Prostatic Neoplasms
- Urinary Bladder Neoplasms

Copyright © 2023 Abnova Corporation. All Rights Reserved.