9F, No. 108, Jhouzih St., Taipei, Taiwan
Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218
E-mail: sales@abnova.com

Datasheet

HDAC1 monoclonal antibody, clone AIB-8

Catalog Number: MAB20120

Regulatory Status: For research use only (RUO)

Product Description: Rabbit monoclonal antibody raised against synthetic peptide of human HDAC1.

Clone Name: AIB-8

Immunogen: A synthetic peptide corresponding to human HDAC1.

Host: Rabbit

Theoretical MW (kDa): 55.103

Reactivity: Human

Applications: ICC, IF, IHC-P, WB-Ce
(See our web site product page for detailed applications information)

Protocols: See our web site at http://www.abnova.com/support/protocols.asp or product page for detailed protocols

Form: Liquid

Purification: Affinity purification

Isotype: IgG

Recommend Usage: Immunocytochemistry (1:50-1:200)
Immunofluorescence (1:50-1:200)
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1:50-1:200)
Western Blot (1:1000-1:2000)
The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS, 150 mM NaCl, pH 7.4 (50% glycerol, 0.02% sodium azide).

Storage Instruction: Store at -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Entrez GeneID: 3065

Gene Symbol: HDAC1

Gene Alias: DKFZp686H12203, GON-10, HD1, RPD3, RPD3L1

Gene Summary: Histone acetylation and deacetylation, catalyzed by multisubunit complexes, play a key role in the regulation of eukaryotic gene expression. The protein encoded by this gene belongs to the histone deacetylase/acuc/apha family and is a component of the histone deacetylase complex. It also interacts with retinoblastoma tumor-suppressor protein and this complex is a key element in the control of cell proliferation and differentiation. Together with metastasis-associated protein-2, it deacetylates p53 and modulates its effect on cell growth and apoptosis. [provided by RefSeq]