


MaxPab@

TRIM23 MaxPab mouse polyclonal antibody (B01P)

Catalog # H00000373-B01P Size 50 ug

Applications



Western Blot (Transfected lysate)

Western Blot analysis of TRIM23 expression in transfected 293T cell line (<u>H00000373-T01</u>) by TRIM23 MaxPab polyclonal antibody.

Lane 1: TRIM23 transfected lysate(63.14 KDa).

Lane 2: Non-transfected lysate.

Specification	
Product Description	Mouse polyclonal antibody raised against a full-length human TRIM23 protein.
Immunogen	TRIM23 (NP_001647, 1 a.a. ~ 574 a.a) full-length human protein.
Sequence	MATLVVNKLGAGVDSGRQGSRGTAVVKVLECGVCEDVFSLQGDKVPRLLLCGHTVCHDCLTRL PLHGRAIRCPFDRQVTDLGDSGVWGLKKNFALLELLERLQNGPIGQYGAAEESIGISGESIIRCDED EAHLASVYCTVCATHLCSECSQVTHSTKTLAKHRRVPLADKPHEKTMCSQHQVHAIEFVCLEEG CQTSPLMCCVCKEYGKHQGHKHSVLEPEANQIRASILDMAHCIRTFTEEISDYSRKLVGIVQHIEGG EQIVEDGIGMAHTEHVPGTAENARSCIRAYFYDLHETLCRQEEMALSVVDAHVREKLIWLRQQQE DMTILLSEVSAACLHCEKTLQQDDCRVVLAKQEITRLLETLQKQQQQFTEVADHIQLDASIPVTFTK DNRVHIGPKMEIRVVTLGLDGAGKTTILFKLKQDEFMQPIPTIGFNVETVEYKNLKFTIWDVGGKHKL RPLWKHYYLNTQAVVFVVDSSHRDRISEAHSELAKLLTEKELRDALLLIFANKQDVAGALSVEEITE LLSLHKLCCGRSWYQGCDARSGMGLYEGLDWLSRQLVAAGVLDVA
Host	Mouse
Reactivity	Human
Quality Control Testing	Antibody reactive against mammalian transfected lysate.
Storage Buffer	In 1x PBS, pH 7.4





Storage Instruction

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

Western Blot (Transfected lysate)

Western Blot analysis of TRIM23 expression in transfected 293T cell line ($\underline{\text{H00000373-T01}}$) by TRIM23 MaxPab polyclonal antibody.

Lane 1: TRIM23 transfected lysate(63.14 KDa).

Lane 2: Non-transfected lysate.

Protocol Download

Gene Info — TRIM23	
Entrez GenelD	<u>373</u>
GeneBank Accession#	NM_001656
Protein Accession#	<u>NP_001647</u>
Gene Name	TRIM23
Gene Alias	ARD1, ARFD1, RNF46
Gene Description	tripartite motif-containing 23
Omim ID	601747
Gene Ontology	<u>Hyperlink</u>
Gene Summary	The protein encoded by this gene is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. This protein is also a member of the ADP ribosylation factor family of guanine nucleotide-binding family of proteins. Its carboxy terminus contains an ADP-ribosylation factor domain and a guanine nucleotide binding site, while the amino terminus contains a GTPase activating protein d omain which acts on the guanine nucleotide binding site. The protein localizes to lysosomes and t he Golgi apparatus. It plays a role in the formation of intracellular transport vesicles, their movement from one compartment to another, and phopholipase D activation. Three alternatively spliced transcript variants for this gene have been described. [provided by RefSeq
Other Designations	ADP-ribosylation factor domain protein 1 ADP-ribosylation factor domain protein 1, 64kDa ARF d omain protein 1 GTP-binding protein ARD-1 OTTHUMP00000123453 tripartite motif protein TRI M23