

## CSTA rabbit monoclonal antibody

Catalog # H00001475-K      Size 100 ug x up to 3

### Specification

<b>Product Description</b>	Rabbit monoclonal antibody raised against a human CSTA peptide using ARM Technology.
<b>Immunogen</b>	A synthetic peptide of human CSTA is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
<b>Host</b>	Rabbit
<b>Library Construction</b>	Non-fusion antibody library from rabbit spleen ( <a href="#">ARM Technology</a> ).
<b>Expression</b>	Overexpression vector and transfection into 293H cell line.
<b>Reactivity</b>	Human
<b>Purification</b>	Protein A
<b>Isotype</b>	IgG
<b>Quality Control Testing</b>	Antibody reactive against human CSTA peptide by ELISA and mammalian transfected lysate by Western Blot.
<b>Storage Buffer</b>	In 1x PBS, pH 7.4
<b>Storage Instruction</b>	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
<b>Deliverable</b>	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
<b>Note</b>	1. Customer may provide cell or tissue lysate for antibody screening. 2. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering including F(ab) <sub>2</sub> , IgG, scFv and different Fc and non-Fc conjugates per customer request.

### Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- ELISA

## Gene Info — CSTA

Entrez GeneID [1475](#)

GeneBank Accession# [CSTA](#)

Gene Name CSTA

Gene Alias STF1, STFA

Gene Description cystatin A (stefin A)

Omim ID [184600](#)

Gene Ontology [Hyperlink](#)

**Gene Summary**

The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibitory activity. There are three inhibitory families in the superfamily, including the type 1 cystatins (stefins), type 2 cystatins, and kininogens. This gene encodes a stefin that functions as a cysteine protease inhibitor, forming tight complexes with papain and the cathepsins B, H, and L. The protein is one of the precursor proteins of cornified cell envelope in keratinocytes and plays a role in epidermal development and maintenance. Stefins have been proposed as prognostic and diagnostic tools for cancer. [provided by RefSeq]

**Other Designations** cystatin A|cystatin AS|stefin A

## Disease

- [Genetic Predisposition to Disease](#)
- [Psoriasis](#)