


RecomAb™

Histone H3 (phospho T6) monoclonal antibody, clone RM160

Catalog # MAB12795 Size 100 ug

Applications



Western Blot (Cell lysate)

Western Blot analysis of Lane 1: acid extracts of HeLa cell treated with Nocodazole and Lane 2: acid extracts of HeLa cell untreated with Histone H3 (phospho T6) monoclonal antibody, clone RM160 (Cat # MAB12795) at 0.1 ug/mL working concentration, showed a band of Histone H3 phosphorylated at Threonine 6.

Specification	
Product Description	Rabbit recombinant monoclonal antibody raised against of human histone H3 (T6).
Antibody Species	Rabbit
lmmunogen	Original antibody is raised against a synthetic phosphopeptide corresponding to residues surroundin g T6 of human Histone H3.
Sequence	N/A
Specificity	This antibody reacts to Histone H3 phosphorylated at Threonine 6. No cross reactivity with other pho sphorylated histones
Form	Liquid
Purification	Protein A purification
Isotype	lgG
Recommend Usage	ELISA (0.2 ug/mL-1 ug/mL) Western Blot (0.1 ug/mL-1 ug/mL) The optimal working dilution should be determined by the end user.



Product Information

Storage Buffer	In PBS (50% glycerol, 1% BSA, 0.09% sodium azide)
Storage Instruction	Store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

Western Blot (Cell lysate)

Western Blot analysis of Lane 1: acid extracts of HeLa cell treated with Nocodazole and Lane 2: acid extracts of HeLa cell untreated with Histone H3 (phospho T6) monoclonal antibody, clone RM160 (Cat # MAB12795) at 0.1 ug/mL working concentration, showed a band of Histone H3 phosphorylated at Threonine 6.

Enzyme-linked Immunoabsorbent Assay

Gene Info — HIST1H3A	
Entrez GenelD	<u>8350</u>
Protein Accession#	<u>P84243</u>
Gene Name	HIST1H3A
Gene Alias	H3/A, H3FA
Gene Description	histone cluster 1, H3a
Omim ID	<u>602810</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chro mosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped ar ound a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H 1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H3 family. Transcripts from this gene lack polyA t ails; instead, they contain a palindromic termination element. This gene is found in the large histon e gene cluster on chromosome 6p22-p21.3. [provided by RefSeq
Other Designations	H3 histone family, member A histone 1, H3a



Pathway

Systemic lupus erythematosus