


CD80 monoclonal antibody, clone AOEH-3

Catalog # MAB19852 Size 100 uL

Applications



Western Blot (Cell lysate)

Western blot analysis of Raji cell lysate with CD80 monoclonal antibody.

Specification

Product Description Rabbit monoclonal antibody raised against synthetic peptide of human CD80.

Immunogen A synthetic peptide corresponding to human CD80.

Host Rabbit

Reactivity Human

Form Liquid

Purification Affinity purification

Isotype IgG

Recommend Usage Immunohistochemistry (1:50-1:200)
Western Blot (1:500-1:2000)
The optimal working dilution should be determined by the end user.

Storage Buffer In PBS, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

Storage Instruction Store at -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Cell lysate)

Western blot analysis of Raji cell lysate with CD80 monoclonal antibody.

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Gene Info — CD80

Entrez GeneID	941
---------------	---------------------

Protein Accession#	P33681
--------------------	------------------------

Gene Name	CD80
-----------	------

Gene Alias	CD28LG, CD28LG1, LAB7
------------	-----------------------

Gene Description	CD80 molecule
------------------	---------------

Omim ID	112203
---------	------------------------

Gene Ontology	Hyperlink
---------------	---------------------------

Gene Summary	The B-lymphocyte activation antigen B7-1 (formerly referred to as B7) provides regulatory signals for T lymphocytes as a consequence of binding to the CD28 (MIM 186760) and CTLA4 (MIM 123890) ligands of T cells.[supplied by OMIM]
--------------	---

Other Designations	B-lymphocyte activation antigen B7 CD80 antigen CD80 antigen (CD28 antigen ligand 1, B7-1 antigen) costimulatory factor CD80 costimulatory molecule variant IgV-CD80
--------------------	--

Pathway

- [Allograft rejection](#)
- [Autoimmune thyroid disease](#)
- [Cell adhesion molecules \(CAMs\)](#)
- [Graft-versus-host disease](#)

- [Systemic lupus erythematosus](#)
- [Toll-like receptor signaling pathway](#)
- [Type I diabetes mellitus](#)

Disease

- [Arthritis](#)
- [Asthma](#)
- [Bronchiolitis](#)
- [Carcinoma](#)
- [Celiac Disease](#)
- [Disease Progression](#)
- [Genetic Predisposition to Disease](#)
- [Hepatitis C](#)
- [Infant](#)
- [Liver Neoplasms](#)
- [Lung Neoplasms](#)
- [Lupus Erythematosus](#)
- [Multiple Sclerosis](#)
- [Papillomavirus Infections](#)
- [Pulmonary Disease](#)
- [Respiratory Syncytial Virus Infections](#)
- [Sarcoidosis](#)
- [Urinary Bladder Neoplasms](#)
- [Uterine Cervical Neoplasms](#)
- [Viremia](#)
- [Werner syndrome](#)