

Bioactive

IL15 (Human) Recombinant Protein

Catalog # P8347

Size 2 x 10 ug

Specification

Product Description	Human IL15 (P40933, 49 a.a. - 162 a.a.) partial recombinant protein with His tag at C-terminus expressed in <i>Escherichia coli</i> .
Sequence	MNWWNVISDLKKIEDLIQSMHIDATLYTESDVHPSCKVTAMKCFLELQVISLESGDASIHDVTENLIILANNSLSSNGNVTESGCKECEELEEKNIKEFLQSFVHIVQMFINTSLEHHHHHH
Host	<i>Escherichia coli</i>
Theoretical MW (kDa)	13.9
Form	Liquid
Preparation Method	<i>Escherichia coli</i> expression system
Purity	> 95% by SDS-PAGE
Activity	ED ₅₀ is < 2.5 ng/mL, measured in a cell proliferation assay using CTLL2 mouse cytotoxic T cells, cell score corresponding to a specific activity of > 4 x 10 ⁵ units/mg.
Recommend Usage	Biological Activity SDS-PAGE The optimal working dilution should be determined by the end user.
Storage Buffer	In 20mM Tris-HCl buffer pH 8.0 (0.2mM PMSF, 10% glycerol)
Storage Instruction	Store at 2°C to 8°C for 1 week. For long term storage, aliquot and store at -20°C to -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Functional Study
- SDS-PAGE

Gene Info — IL15

Entrez GeneID [3600](#)

Protein Accession# [P40933](#)

Gene Name IL15

Gene Alias IL-15, MGC9721

Gene Description interleukin 15

Omim ID [600554](#)

Gene Ontology [Hyperlink](#)

Gene Summary

The protein encoded by this gene is a cytokine that regulates T and natural killer cell activation and proliferation. This cytokine and interleukine 2 share many biological activities. They are found to bind common hematopoietin receptor subunits, and may compete for the same receptor, and thus negatively regulate each other's activity. The number of CD8+ memory cells is shown to be controlled by a balance between this cytokine and IL2. This cytokine induces the activation of JAK kinases, as well as the phosphorylation and activation of transcription activators STAT3, STAT5, and STAT6. Studies of the mouse counterpart suggested that this cytokine may increase the expression of apoptosis inhibitor BCL2L1/BCL-x(L), possibly through the transcription activation activity of STAT6, and thus prevent apoptosis. Two alternatively spliced transcript variants of this gene encoding the same protein have been reported. [provided by RefSeq]

Other Designations OTTHUMP00000164617

Pathway

- [Cytokine-cytokine receptor interaction](#)
- [Jak-STAT signaling pathway](#)

Disease

- [Adenocarcinoma](#)
- [Arthritis](#)
- [Asthma](#)
- [Bronchial Hyperreactivity](#)
- [Bronchiolitis](#)

- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Celiac Disease](#)
- [Connective Tissue Diseases](#)
- [Coronary Disease](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Esophageal Neoplasms](#)
- [Fetal Diseases](#)
- [Fractures](#)
- [Genetic Predisposition to Disease](#)
- [Hematologic Diseases](#)
- [Hodgkin Disease](#)
- [Hypersensitivity](#)
- [Hypertrophy](#)
- [Infant](#)
- [Infection](#)
- [Inflammation](#)
- [Kidney Failure](#)
- [Lung Neoplasms](#)
- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Metabolic Syndrome X](#)
- [Multiple Myeloma](#)
- [Musculoskeletal Diseases](#)
- [Neoplasm](#)

- [Obesity](#)
- [Occupational Diseases](#)
- [Osteoporosis](#)
- [Pregnancy Complications](#)
- [Premature Birth](#)
- [Psoriasis](#)
- [Pulmonary Disease](#)
- [Recurrence](#)
- [Respiratory Syncytial Virus Infections](#)
- [Skin Diseases](#)
- [Urinary Bladder Neoplasms](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)