

Goat Anti-Dog IgG (H&L) secondary antibody (Fluorescein)

Catalog # PAB10523

Size 2 mg

Specification

Product Description	Goat anti-dog IgG recognizes dog IgG whole molecule. This secondary antibody was purified using a ntigen affinity chromatography. The antibody is conjugated with Fluorescein.
Immunogen	Dog IgG whole molecule
Host	Goat
Reactivity	Dog
Form	Liquid
Conjugation	FITC
Purification	This product was prepared from monospecific antiserum by immunoaffinity chromatography, followed by solid phase adsorption(s) to remove any unwanted reactivities.
Conjugation Note	3.6 moles DTAF per mole of goat IgG
Recommend Usage	The optimal working dilution should be determined by the end user.
Storage Buffer	In 20 mM KH ₂ PO ₄ , 150 mM NaCl, pH 7.2 (0.01% sodium azide)
Storage Instruction	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunofluorescence
- Flow Cytometry

Publication Reference

- [Immunofluorescence using dichlorotriazinylaminofluorescein \(DTAF\). I. Preparation and fractionation of labelled IgG.](#)

D Blakeslee, M G Baines.

Journal of Immunological Methods 1976 Dec; 13(3-4):305.