

# Goat Anti-Golden Syrian hamster IgG (H&L) secondary antibody (Rhodamine)

Catalog # PAB10592

Size 2 mg

## Specification

<b>Product Description</b>	Goat anti-Golden Syrian hamster IgG recognizes Golden Syrian hamster IgG whole molecule. This secondary antibody was purified using antigen affinity chromatography. The antibody is conjugated with Rhodamine.
<b>Immunogen</b>	Golden Syrian hamster IgG whole molecule
<b>Host</b>	Goat
<b>Reactivity</b>	Hamster
<b>Form</b>	Lyophilized
<b>Conjugation</b>	Rhodamine (TRITC)
<b>Purification</b>	This product was prepared from monospecific antiserum by immunoaffinity chromatography, followed by solid phase adsorption(s) to remove any unwanted reactivities.
<b>Conjugation Note</b>	2.8 moles TRITC per mole of goat IgG
<b>Recommend Usage</b>	The optimal working dilution should be determined by the end user.
<b>Storage Buffer</b>	Lyophilized from 20 mM $\text{KH}_2\text{PO}_4$ , 150 mM NaCl, pH 7.2 (0.01% sodium azide)
<b>Storage Instruction</b>	Store at 4°C on dry atmosphere. After reconstitution with 1 mL of deionized water, store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
<b>Note</b>	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

## Applications

- Immunofluorescence

- Flow Cytometry

## Publication Reference

- [Texas Red, a hydrophilic, red-emitting fluorophore for use with fluorescein in dual parameter flow microfluorometric and fluorescence microscopic studies.](#)

J A Titus, R Haugland, S O Sharrow, D M Segal.

Journal of Immunological Methods 1982 Apr; 50(2):193.