

Goat Anti-Pig IgG (H&L) secondary antibody (Fluorescein)

Catalog # PAB10787

Size 2 mg

Specification

Product Description	Goat anti-pig IgG recognizes pig IgG whole molecule. This secondary antibody was purified using antigen affinity chromatography. The antibody is conjugated with Fluorescein.
Immunogen	Pig IgG whole molecule
Host	Goat
Reactivity	Pig
Form	Lyophilized
Conjugation	FITC
Purification	This product was prepared from monospecific antiserum by immunoaffinity chromatography, followed by solid phase adsorption(s) to remove any unwanted reactivities.
Conjugation Note	3.1 moles FITC per mole of goat IgG
Recommend Usage	The optimal working dilution should be determined by the end user.
Storage Buffer	Lyophilized from 10 mM potassium phosphate, 75 mM NaCl, 50% (v/v) glycerol, pH 7.2 (0.01% (w/v) sodium azide, 10 mg/mL BSA (Immunoglobulin Lyophilized from and Protease free))
Storage Instruction	Store at 4°C on dry atmosphere. After reconstitution with 1 mL of deionized water, store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunofluorescence
- Flow Cytometry

Publication Reference

- [Conjugation of fluorescein isothiocyanate to antibodies. I. Experiments on the conditions of conjugation.](#)

The TH, Feltkamp TE.

Immunology 1970 Jun; 18(6):865.

Application: Conjugation, Fluorescein isothiocyanate (FITC)