


IL15 polyclonal antibody

Catalog # PAB27236 Size 100 ug

Applications



Western Blot (Tissue lysate)

Western blot analysis of IL15 in rat spleen tissue with IL15 polyclonal antibody (Cat # PAB27236) at (lane 1) 1 and (lane 2) 2 ug/mL.

Specification

Product Description	Rabbit polyclonal antibody raised against synthetic peptide of IL15.
Immunogen	A synthetic peptide corresponding to 13 amino acids at internal region of human IL15.
Host	Rabbit
Reactivity	Human, Mouse, Rat
Specificity	At least two isoforms of IL-15 are known to exist.
Form	Liquid
Purification	Peptide affinity purification
Concentration	1 mg/mL
Isotype	IgG
Recommend Usage	Western Blot (1-2 ug/mL) Immunohistochemistry (2.5 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.02% sodium azide)

Storage Instruction

Store at 4°C for three months. For long term storage store at -20°C.
Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Tissue lysate)

Western blot analysis of IL15 in rat spleen tissue with IL15 polyclonal antibody (Cat # PAB27236) at (lane 1) 1 and (lane 2) 2 ug/mL.

- Immunohistochemistry

- Enzyme-linked Immunoabsorbent Assay

Gene Info — IL15

Entrez GeneID
[3600](#)
Protein Accession#
[NP_000576](#)
Gene Name

IL15

Gene Alias

IL-15, MGC9721

Gene Description

interleukin 15

Omim ID
[600554](#)
Gene Ontology
[Hyperlink](#)
Gene Summary

The protein encoded by this gene is a cytokine that regulates T and natural killer cell activation and proliferation. This cytokine and interleukine 2 share many biological activities. They are found to bind common hematopoietin receptor subunits, and may compete for the same receptor, and thus negatively regulate each other's activity. The number of CD8+ memory cells is shown to be controlled by a balance between this cytokine and IL2. This cytokine induces the activation of JAK kinases, as well as the phosphorylation and activation of transcription activators STAT3, STAT5, and STAT6. Studies of the mouse counterpart suggested that this cytokine may increase the expression of apoptosis inhibitor BCL2L1/BCL-x(L), possibly through the transcription activation activity of STAT6, and thus prevent apoptosis. Two alternatively spliced transcript variants of this gene encoding the same protein have been reported. [provided by RefSeq]

Pathway

- [Cytokine-cytokine receptor interaction](#)
- [Jak-STAT signaling pathway](#)

Disease

- [Adenocarcinoma](#)
- [Arthritis](#)
- [Asthma](#)
- [Bronchial Hyperreactivity](#)
- [Bronchiolitis](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Celiac Disease](#)
- [Connective Tissue Diseases](#)
- [Coronary Disease](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Esophageal Neoplasms](#)
- [Fetal Diseases](#)
- [Fractures](#)
- [Genetic Predisposition to Disease](#)
- [Hematologic Diseases](#)
- [Hodgkin Disease](#)
- [Hypersensitivity](#)

- [Hypertrophy](#)
- [Infant](#)
- [Infection](#)
- [Inflammation](#)
- [Kidney Failure](#)
- [Lung Neoplasms](#)
- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Metabolic Syndrome X](#)
- [Multiple Myeloma](#)
- [Musculoskeletal Diseases](#)
- [Neoplasm](#)
- [Obesity](#)
- [Occupational Diseases](#)
- [Osteoporosis](#)
- [Pregnancy Complications](#)
- [Premature Birth](#)
- [Psoriasis](#)
- [Pulmonary Disease](#)
- [Recurrence](#)
- [Respiratory Syncytial Virus Infections](#)
- [Skin Diseases](#)
- [Urinary Bladder Neoplasms](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)