

# ACPP polyclonal antibody

Catalog # PAB30373 Size 100 uL

## **Applications**



# Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human prostate with ACPP polyclonal antibody (Cat # PAB30373) shows strong cytoplasmic positivity in glandular cells at 1:200-1:500 dilution.

Specification	
Product Description	Rabbit polyclonal antibody raised against partial recombinant human ACPP.
Immunogen	Recombinant protein corresponding to human ACPP.
Sequence	QLLYLPFRNCPRFQELESETLKSEEFQKRLHPYKDFIATLGKLSGLHGQDLFGIWSKVYDPLYCES VHNFTLPSWATEDTMTKLRELSELSLLSLYGIHKQKEKSRLQGGVLVNEILNHMKRAT
Host	Rabbit
Reactivity	Human
Form	Liquid
Purification	Antigen affinity purification
Isotype	lgG
Recommend Usage	Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1:200-1:500) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH 7.2 (40% glycerol, 0.02% sodium azide).



#### **Product Information**

Storage Instruction	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

## **Applications**

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human prostate with ACPP polyclonal antibody (Cat # PAB30373) shows strong cytoplasmic positivity in glandular cells at 1:200-1:500 dilution.

Gene Info — ACPP	
Entrez GenelD	<u>55</u>
Protein Accession#	<u>P15309</u>
Gene Name	ACPP
Gene Alias	ACP-3, ACP3, PAP
Gene Description	acid phosphatase, prostate
Omim ID	<u>171790</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	This gene encodes an enzyme that catalyzes the conversion of orthophosphoric monoester to alc ohol and orthophosphate. It is synthesized under androgen regulation and is secreted by the epith elial cells of the prostate gland. An alternatively spliced transcript variant encoding a longer isofor m has been found for this gene. This isoform contains a transmembrane domain and is localized in the plasma membrane-endosomal-lysosomal pathway. [provided by RefSeq
Other Designations	prostatic acid phosphotase

## Pathway

- gamma-Hexachlorocyclohexane degradation
- Riboflavin metabolism



### Disease

Tobacco Use Disorder