


RecomAb™

BAD recombinant monoclonal antibody, clone BADS112-B9

Catalog # RAB02822 Size 200 uL

Applications



Flow Cytometry

Peptide blocking flow cytometric analysis of U937 cells secondary antibody only negative control (blue) or Calyculin A-treated (green) or CalA and blocked with phospho peptide (gold) or CalA and blocked with non-phospho peptide (purple) using Phospho-BAD (Ser112) antibody BADS112-B9 at 0.1 ug/mL.

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human BAD.
Antibody Species	Rabbit
Immunogen	A synthetic phospho-peptide corresponding to residues surrounding Ser112 of human phospho BAD
Reactivity	Human
Form	Liquid
Purification	Protein A+G
Isotype	Rabbit IgG1L
Recommend Usage	Flow Cytometry The optimal working dilution should be determined by the end user.
Storage Buffer	1X PBS, 0.02% Sodium azide, 50% Glycerol, 0.1% BSA

Storage Instruction

Store at -20°C.
Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Flow Cytometry

Peptide blocking flow cytometric analysis of U937 cells secondary antibody only negative control (blue) or Calyculin A-treated (green) or CalA and blocked with phospho peptide (gold) or CalA and blocked with non-phospho peptide (purple) using Phospho-BAD (Ser112) antibody BADS112-B9 at 0.1 ug/mL.

Gene Info — BAD

Entrez GeneID
[572](#)
Protein Accession#
[Q92934](#)
Gene Name

BAD

Gene Alias

BBC2, BCL2L8

Gene Description

BCL2-associated agonist of cell death

Omim ID
[603167](#)
Gene Ontology
[Hyperlink](#)
Gene Summary

The protein encoded by this gene is a member of the BCL-2 family. BCL-2 family members are known to be regulators of programmed cell death. This protein positively regulates cell apoptosis by forming heterodimers with BCL-xL and BCL-2, and reversing their death repressor activity. Proapoptotic activity of this protein is regulated through its phosphorylation. Protein kinases AKT and MAP kinase, as well as protein phosphatase calcineurin were found to be involved in the regulation of this protein. Alternative splicing of this gene results in two transcript variants which encode the same isoform. [provided by RefSeq]

Other Designations

BCL-X/BCL-2 binding protein|BCL2-antagonist of cell death protein|BCL2-binding component 6|BCL2-binding protein

Pathway

- [Acute myeloid leukemia](#)

- [Amyotrophic lateral sclerosis \(ALS\)](#)
- [Apoptosis](#)
- [Chronic myeloid leukemia](#)
- [Colorectal cancer](#)
- [Endometrial cancer](#)
- [ErbB signaling pathway](#)
- [Focal adhesion](#)
- [Insulin signaling pathway](#)
- [Melanoma](#)
- [Neurotrophin signaling pathway](#)
- [Non-small cell lung cancer](#)
- [Pancreatic cancer](#)
- [Pathways in cancer](#)
- [Prostate cancer](#)
- [VEGF signaling pathway](#)

Disease

- [Cardiovascular Diseases](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Lymphoma](#)
- [Parkinson disease](#)
- [Thyroid Neoplasms](#)