


RecomAb™

CD80 recombinant monoclonal antibody, clone 7C10

Catalog # RAB03341 Size 200 ug

Applications



Immunofluorescence

Immunofluorescent staining of Daudi cells with CD80 recombinant monoclonal antibody, clone 7C10 (Cat # RAB03341).

Immunofluorescence analysis of paraformaldehyde fixed Daudi cells on Shi-fix™ coverslips stained with the chimeric r version of RAB03341 at 10 ug/ml for 1h followed by Alexa Fluor® 488 secondary antibody (2 ug/ml)- showing membrane staining. The nuclear stain is DAPI (blue). Panels show from left-right- top-bottom RAB03341- DAPI- merged channels and an isotype control. The isotype control was an unknown specificity antibody (3.0) followed by staining with Alexa Fluor® 488 secondary antibody.

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human CD80.
Antibody Species	Rabbit
Immunogen	Original antibody is raised against human B7.1 antigen.
Reactivity	Human
Form	Liquid
Isotype	IgG lambda
Recommend Usage	Blocking ELISA Immunofluorescence The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.02% Proclin 300)

Storage Instruction

Store at 4°C for 3 months. For long term storage store at -20°C.
Aliquot to avoid repeated freezing and thawing.

Applications

- Immunofluorescence

Immunofluorescent staining of Daudi cells with CD80 recombinant monoclonal antibody, clone 7C10 (Cat # RAB03341). Immunofluorescence analysis of paraformaldehyde fixed Daudi cells on Shi-fix™ coverslips stained with the chimeric r version of RAB03341 at 10 ug/ml for 1h followed by Alexa Fluor® 488 secondary antibody (2 ug/ml)- showing membrane staining. The nuclear stain is DAPI (blue). Panels show from left-right- top-bottom RAB03341- DAPI- merged channels and an isotype control. The isotype control was an unknown specificity antibody (3.0) followed by staining with Alexa Fluor® 488 secondary antibody.

- Enzyme-linked Immunoabsorbent Assay

- Blocking

Gene Info — CD80

Entrez GeneID	941
Gene Name	CD80
Gene Alias	CD28LG, CD28LG1, LAB7
Gene Description	CD80 molecule
Omim ID	112203
Gene Ontology	Hyperlink
Gene Summary	The B-lymphocyte activation antigen B7-1 (formerly referred to as B7) provides regulatory signals for T lymphocytes as a consequence of binding to the CD28 (MIM 186760) and CTLA4 (MIM 123 890) ligands of T cells.[supplied by OMIM]
Other Designations	B-lymphocyte activation antigen B7 CD80 antigen CD80 antigen (CD28 antigen ligand 1, B7-1 antigen) costimulatory factor CD80 costimulatory molecule variant IgV-CD80

Pathway

- [Allograft rejection](#)
- [Autoimmune thyroid disease](#)

- [Cell adhesion molecules \(CAMs\)](#)
- [Graft-versus-host disease](#)
- [Systemic lupus erythematosus](#)
- [Toll-like receptor signaling pathway](#)
- [Type I diabetes mellitus](#)

Disease

- [Arthritis](#)
- [Asthma](#)
- [Bronchiolitis](#)
- [Carcinoma](#)
- [Celiac Disease](#)
- [Disease Progression](#)
- [Genetic Predisposition to Disease](#)
- [Hepatitis C](#)
- [Infant](#)
- [Liver Neoplasms](#)
- [Lung Neoplasms](#)
- [Lupus Erythematosus](#)
- [Multiple Sclerosis](#)
- [Papillomavirus Infections](#)
- [Pulmonary Disease](#)
- [Respiratory Syncytial Virus Infections](#)
- [Sarcoidosis](#)
- [Urinary Bladder Neoplasms](#)
- [Uterine Cervical Neoplasms](#)

- [Viremia](#)
- [Werner syndrome](#)